

Clinical Cancer Research



A Novel Kinase Inhibitor, INCB28060, Blocks c-MET–Dependent Signaling, Neoplastic Activities, and Cross-Talk with EGFR and HER-3

Xiangdong Liu, Qian Wang, Gengjie Yang, et al.

Clin Cancer Res 2011;17:7127-7138. Published OnlineFirst September 14, 2011.

Updated version	Access the most recent version of this article at: doi: 10.1158/1078-0432.CCR-11-1157
Supplementary Material	Access the most recent supplemental material at: http://clincancerres.aacrjournals.org/content/suppl/2011/09/14/1078-0432.CCR-11-1157.DC1.html

Cited Articles	This article cites by 31 articles, 15 of which you can access for free at: http://clincancerres.aacrjournals.org/content/17/22/7127.full.html#ref-list-1
Citing articles	This article has been cited by 3 HighWire-hosted articles. Access the articles at: http://clincancerres.aacrjournals.org/content/17/22/7127.full.html#related-urls

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
Permissions	To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org .

A Novel Kinase Inhibitor, INCB28060, Blocks c-MET–Dependent Signaling, Neoplastic Activities, and Cross-Talk with EGFR and HER-3

Xiangdong Liu, Qian Wang, Gengjie Yang, Cindy Marando, Holly K. Koblish, Leslie M. Hall, Jordan S. Fridman, Elham Behshad, Richard Wynn, Yu Li, Jason Boer, Sharon Diamond, Chunhong He, Meizhong Xu, Jincong Zhuo, Wenqing Yao, Robert C. Newton, and Peggy A. Scherle

Abstract

Purpose: The c-MET receptor tyrosine kinase plays important roles in the formation, progression, and dissemination of human cancer and presents an attractive therapeutic target. This study describes the preclinical characterization of INCB28060, a novel inhibitor of c-MET kinase.

Experimental Design: Studies were conducted using a series of *in vitro* and *in vivo* biochemical and biological experiments.

Results: INCB28060 exhibits picomolar enzymatic potency and is highly specific for c-MET with more than 10,000-fold selectivity over a large panel of human kinases. This inhibitor potently blocks c-MET phosphorylation and activation of its key downstream effectors in c-MET–dependent tumor cell lines. As a result, INCB28060 potently inhibits c-MET–dependent tumor cell proliferation and migration and effectively induces apoptosis *in vitro*. Oral dosing of INCB28060 results in time- and dose-dependent inhibition of c-MET phosphorylation and tumor growth in c-MET–driven mouse tumor models, and the inhibitor is well tolerated at doses that achieve complete tumor inhibition. In a further exploration of potential interactions between c-MET and other signaling pathways, we found that activated c-MET positively regulates the activity of epidermal growth factor receptors (EGFR) and HER-3, as well as expression of their ligands. These effects are reversed with INCB28060 treatment. Finally, we confirmed that circulating hepatocyte growth factor levels are significantly elevated in patients with various cancers.

Conclusions: Activated c-MET has pleiotropic effects on multiple cancer-promoting signaling pathways and may play a critical role in driving tumor cell growth and survival. INCB28060 is a potent and selective c-MET kinase inhibitor that may have therapeutic potential in cancer treatment. *Clin Cancer Res*; 17(22): 7127–38. ©2011 AACR.

Introduction

c-MET, originally identified as a *TRP–MET* fusion gene from a human osteosarcoma cell line, encodes a prototype member of a distinct subfamily of heterodimeric receptor tyrosine kinases (RTK; ref. 1). Hepatocyte growth factor (HGF) is the only known high affinity ligand for the c-MET receptor. Binding of HGF to c-MET causes receptor multimerization, phosphorylation, and catalytic activation. The activated receptor subsequently recruits adaptor proteins like GAB1 [growth factor receptor–binding protein 2

(GRB2)-associated binding protein], GRB2, SHC (Src homology and collagen), and c-Cbl and leads to activation of multiple downstream effector pathways or proteins, including RAS/MAPK (mitogen-activated protein kinase), PI3K (phosphoinositide-3-kinase)/AKT, FAK (focal adhesion kinase), STAT3/5, RAC/RHO, PLC- γ (phospholipase C γ), c-SRC, SHP2 (a Src homology 2-containing tyrosine phosphatase), and CRKL, that are essential for regulating cell growth, survival, motility, invasion, and cytoskeletal changes (1, 2).

Although c-MET and HGF are present in many different tissues, their expression is mainly confined to cells of epithelial and mesenchymal origins, respectively. The normal functions of the c-MET pathway are largely restricted to organ morphogenesis during development as well as to tissue damage repair and regeneration in adults (1–7). Importantly, aberrant c-MET activation has been frequently found in many human solid tumors and hematologic malignancies (1, 2, 8). The evidence that links c-MET directly to cancer arose from the original discovery of missense *c-MET* mutations in all patients with hereditary

Author's Affiliation: Incyte Corporation, Wilmington, Delaware

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org>).

Corresponding Author: Xiangdong Liu, Incyte Corporation, Route 141 and Henry Clay Road, Building E400/3229C, Wilmington, DE 19880. Phone: 302-498-6815; Fax: 302-425-2760; E-mail: xliu@incyte.com

doi: 10.1158/1078-0432.CCR-11-1157

©2011 American Association for Cancer Research.

Translational Relevance

The c-MET pathway plays important roles in human cancer and presents an attractive therapeutic target. This study describes preclinical characterization of INCB28060, a selective and potent c-MET kinase inhibitor currently being studied in patients. INCB28060 effectively inhibits c-MET-dependent signaling and neoplastic activities in cell and animal models. In exploring potential cross-talk between c-MET and EGFR/HER-3, we find that c-MET upregulates the activity of EGFR and HER-3 directly through these receptors or by stimulating their ligand production, and these effects can be reversed by INCB28060. Although c-MET pathway inhibitors are being developed, they have limitations in their potency, selectivity, pharmacokinetic property, safety profile, and/or mechanism of action, which hampered their use in assessing the true therapeutic potential of c-MET inhibition in patients. The properties displayed by INCB28060 may enable this agent to overcome the limitations of other c-MET pathway inhibitors and specifically address the importance of this pathway in human cancers.

papillary renal cell carcinoma (PRCC) and a small subset (13%) of sporadic PRCC samples studied (9). Trisomy of chromosome 7, where both *HGF* and *c-MET* genes are located, also occurs frequently in patients with PRCC. Subsequently, *c-MET* mutations were found in many other cancers, including gastric, head and neck, liver, ovarian, non-small cell lung (NSCL), and thyroid cancers, as well as in the metastases of some of these cancers (2, 8, 10–12). The tumorigenic potential of several of the reported c-MET mutants was confirmed in preclinical models (13, 14). In addition to mutation, c-MET can also be activated via receptor overexpression and amplification, or by elevated HGF levels. Amplification and/or overexpression of *c-MET* have been reported in patients with various cancer types, including brain, colorectal, gastric, lung, head, and neck cancers, and elevated HGF levels have been found in most human cancers (2, 8, 15–17). Furthermore, published data suggest that both c-MET and HGF elevations have been associated with poor clinical outcomes (8, 17–21), underscoring the importance of increased c-MET signaling in human cancers. Finally, c-MET signaling may also be modulated through its interactions with other cell surface proteins, ranging from integrins to members of the epithelial growth factor receptor (EGFR) tyrosine kinase family, suggesting that a network of interacting proteins may modulate and eventually determine the magnitude and duration of c-MET signaling in a cell context-dependent manner (10).

At the cellular level, activated c-MET promotes tumor cell growth, survival, migration, and invasion, as well as tumor angiogenesis (1, 2). Previous studies have validated the c-MET pathway as an attractive target for cancer intervention

(2). First, overexpression of HGF and/or c-MET (wild-type or mutants) conferred a transforming phenotype in cell lines and mice expressing *HGF* and/or *c-MET* as a transgene(s) developed different types of tumors and metastatic lesions. Second, downregulation of HGF or c-MET expression or inhibition of c-MET kinase activity in HGF/c-MET-driven tumor cells significantly decreased cell growth, survival, motility, migration and invasion *in vitro*, and reduced tumorigenic and metastatic potential as well as tumor angiogenesis *in vivo*. Recent data from studies of anti-c-MET or anti-HGF antibodies and small molecule c-MET tyrosine kinase inhibitors (TKI) further support the therapeutic potential of c-MET inhibition (10).

Despite the strong evidence supporting the critical role of the c-MET pathway in many human cancers, no c-MET inhibitors or c-MET pathway antagonists have been approved for treating cancer patients. Some of the c-MET pathway-targeted agents currently being evaluated in clinical trials have limitations in terms of their potency, selectivity, pharmacokinetic property, safety profile, and/or specific mechanism of action, which may significantly hamper their ability to achieve optimal inhibition of the pathway at a better tolerated dose in cancer patients (10). Here we describe the identification and characterization of INCB28060, a novel orally active inhibitor of c-MET kinase with properties that may overcome the aforementioned limitations. INCB28060 potently and specifically inhibits c-MET enzyme activity, c-MET-mediated signal transduction, and the c-MET-dependent neoplastic phenotype of tumor cells *in vitro*. The compound exhibits strong antitumor activity in c-MET-dependent tumor models at well-tolerated doses. Furthermore, we found that activated c-MET upregulates the cancer-promoting EGFR and HER-3 pathways. Therefore, inhibition of increased c-MET activity using INCB28060 may have therapeutic potential in human cancer and this is currently being explored in phase I clinical trials.

Materials and Methods

Cell lines, human samples, and reagents

All human cancer cell lines (SNU-5, SNU-1, U-87MG, 786-O, A549, H441, H596, H1437, H1993, BT474, A549, and HT-29) were purchased from the American Type Culture Collection (ATCC) and routinely maintained according to ATCC's recommendations. MKN-45 cell line was obtained from German Collection of Microorganisms and Cell Cultures and maintained according to supplier's recommendation. The S114 cell line that stably expresses human HGF and c-MET was obtained from the NIH (Rockville, MD). Blood samples were obtained from healthy volunteers and cancer patients after informed consent was obtained in accordance with the Declaration of Helsinki and Institutional Review Board approval was received from Incyte. INCB28060 was synthesized at Incyte Corporation as described previously (22). The compound was prepared as a 5 mmol/L stock solution in 100% dimethyl sulfoxide and routinely stored at room

temperature. Actinomycin D and cycloheximide were purchased from Sigma-Aldrich.

In vitro c-MET kinase assay

The assay was conducted and compound potency was determined as described previously (23).

Western blotting

To prepare whole-cell protein extracts, cells maintained in appropriate media were treated for 2 hours with INCB28060 at concentrations indicated in individual experiments. If HGF treatment was required, cells were usually starved in appropriate media containing 0.2% FBS for at least 24 hours and followed by pretreatment with INCB28060 for 2 hours. Cells were then stimulated with 50 ng/mL recombinant human HGF (R&D Systems) for an additional 15 minutes. Whole-cell protein extracts were prepared and Western blot analysis was carried out as described previously (24). All antibodies were purchased from commercial sources. Antibodies obtained from Cell Signaling Technology include rabbit anti-phospho-c-MET (Tyr1234/1235) mAb, rabbit anti-phospho-c-MET (Tyr1003), mouse anti-c-MET mAb, rabbit anti-phospho-STAT3 (Tyr705) mAb, anti-phospho-STAT5 (Tyr694) mAb, rabbit anti-phospho-EGFR (Tyr1173) mAb, rabbit anti-EGFR antibody, rabbit anti-phospho-HER-3/ErbB3 (Tyr1289) mAb, rabbit anti-phospho-p44/42 MAP kinase (Thr202/Tyr204) polyclonal Ab, rabbit anti-p44/42 MAPK [extracellular signal-regulated kinase (Erk1/2)] polyclonal Ab, rabbit anti-phospho-AKT (Ser473) Ab, rabbit anti-AKT polyclonal Ab, rabbit anti-phospho-GAB1 (Tyr627) polyclonal Ab, rabbit anti-phospho-FAK (Tyr576/577) polyclonal Ab, HRP-linked anti-rabbit IgG Ab and HRP-linked anti-mouse IgG Ab. Antibodies from Santa Cruz Biotechnology include rabbit anti-human c-MET polyclonal Ab, rabbit anti-phospho-c-MET (Tyr1349), rabbit anti-phospho-ROK β polyclonal Ab and rabbit anti-ROK polyclonal Ab. Mouse anti-ErbB-3/HER-3 mAb was obtained from Millipore. Recombinant human MSP (macrophage stimulating protein) was purchased from R&D Systems. Western blot chemiluminescence detection reagents were obtained from Pierce. All antibodies and related agents were stored and used according to manufacturers' instructions.

Immunoprecipitation

The entire procedure was carried out at 4°C. Approximately 200 to 300 μ g of whole-cell protein extracts in 400 μ L protein lysis buffer was mixed and incubated with anti-c-MET primary antibody for 1 hour. Twenty to 40 μ L of resuspended volume of protein A/G agarose (Santa Cruz Biotechnology) was added to the mixture and incubated on a rotating device overnight. The pellet was collected by centrifugation at 2,500 rpm for 5 minutes and the supernatant was aspirated and discarded. The pellet was washed 4 times with lysis buffer and after the final wash, the supernatant was aspirated and discarded, and the pellet was resuspended in 20 μ L of 1 \times electrophoresis sample buffer. The samples were boiled for 2 minutes prior to electrophoresis. Western blot analysis was carried out as described above.

ELISAs

ELISA-based detections of phospho-c-MET in cell protein extracts, HGF in patient plasma, and TGF- α , AR, and HGR- β 1 in supernatants of cultured cells were obtained using respective ELISA kits from R&D systems according to the manufacturer's instructions. Before analyzing the samples, optimal amounts of protein extracts, human plasma or culture supernatants were determined for individual assays. The ELISA procedures were carried out as following: various amounts of protein extracts, human plasma or culture supernatants were captured with a primary antibody specific for the individual protein to be measured for 2 hours in a 96-well microplate. After washes, an HRP-conjugated detection antibody (an anti-phospho-tyrosine antibody was used for phospho-c-MET detection) was added and incubated for 2 hours. After additional washes, 100 μ L substrate solution was added into each well and the reaction was stopped with 2 N H₂SO₄ within an appropriate amount of time during color development. The optical density was measured in the linear range using a microplate reader at 450 nm with wavelength correction at 540 nm. IC₅₀ values were calculated using GraphPad Prism software.

Cell viability assay

Optimal cell density used in the viability assay was predetermined for individual cell lines. To determine compound potency, cells were seeded into 96-well microplates at the appropriate density in media containing 1% to 2% FBS and supplemented with serial dilutions of INCB28060 in a final volume of 100 μ L per well. After 72-hour incubation, 24 μ L of CellTiter 96 Aqueous One Solution (Promega) was added to each well, and the plates were incubated for 2 hours in a 37°C incubator. The optical density was measured in the linear range using a microplate reader at 490 nm with wavelength correction at 650 nm. IC₅₀ values were calculated using the GraphPad Prism Software.

Soft agar colony formation assay

U-87MG or H441 cells were prepared at adequate densities in 6-well plates mixed with 0.5 mL top layer agar containing 0.3% agarose (Cambrex) in appropriate culture medium and supplemented with 1% or 10% FBS, in the presence or absence of 50 ng/mL recombinant human HGF and INCB28060 at various concentrations. Cells were evenly laid over 1 mL solidified base layer agar containing 0.6% agarose in culture medium. The plates were incubated at 37°C in a humidified incubator supplied with 5% CO₂. Cells were fed once a week with top agar containing appropriate concentrations of human HGF and INCB28060. The number and size of colonies were evaluated 2 to 3 weeks later when representative photographs were taken.

Cell migration assay

H441 cells were seeded in RPMI-1640 medium containing 10% FBS and grown to complete confluence. Gaps were introduced by scraping cells with a P200 pipette tip. Cells were then stimulated with 50 ng/mL recombinant human

HGF to induce migration across the gap in the presence of various concentrations of INCB28060. After an overnight incubation, representative photographs were taken and a semiquantitative assessment of inhibition of cell migration was conducted.

Apoptosis assay

Cells were seeded in a 96-well plate and grown overnight in culture medium containing 0.5% FBS. Cells were then treated with INCB28060 at various concentrations for 24 hours. Apoptosis was measured using a DNA fragmentation–based Cell Death Detection ELISA^{Plus} kit (Roche Applied Science) according to the manufacturer's instructions. To measure PARP cleavage, cells were grown in 10 cm dishes and treated similarly with INCB28060 as described above. Protein extracts were then prepared and subjected to Western blot analysis using a rabbit anti-cleaved PARP (Asp214) antibody (Cell Signaling Technology).

Animal studies

Animals were housed in barrier facilities fully accredited by the Association for Assessment and Accreditation of Laboratory Animal Care, International. All of the procedures were conducted in accordance with the U.S. Public Health Service Policy on Humane Care and Use of Laboratory Animals and with Incyte Animal Care and Use Committee guidelines. Eight-week-old female Balb/c *nu/nu* mice (Charles River) were inoculated subcutaneously with 4×10^6 tumor cells (S114 model) or with 5×10^6 tumor cells (U-87MG glioblastoma model). Tumors were measured after becoming visible 2 or 3 times weekly in 2 dimensions using a caliper, and the volumes calculated in cubic millimeter using the formula: $V = 0.5 (A \times B^2)$, where *A* and *B* are the long and short diameters of the tumor, respectively. Tumor-bearing animals were sorted into groups with similar mean tumor volumes prior to treatment—usually 50 to 100 mm³ for S114 and 100 to 200 mm³ for U-87MG. Treatments are listed in each experiment.

Efficacy studies

Tumor-bearing mice were dosed orally, twice each day with 1, 3, 10, or 30 mg/kg of free base INCB28060 reconstituted in 5% DMAC in 0.5% methylcellulose for up to 2 weeks. Body weights were monitored throughout the study as a gross measure of toxicity/morbidity. Tumor growth inhibition, expressed in percent, was calculated using the formula: $(1 - [(volume (treated))/volume (vehicle)]) \times 100$.

Pharmacodynamic analysis

For pharmacodynamic analysis, S114 tumor-bearing mice were monitored for tumor growth and then randomized into groups of 3 with average tumor sizes of approximately 300 to 500 mm³. For time course studies, mice were given a single oral dose of 3 mg/kg INCB28060 reconstituted in 5% DMAC in 0.5% methylcellulose and tumors were harvested at the indicated time points. For dose escalation studies, mice were given a single oral dose of INCB28060 at 0.03, 0.1, 0.3, 1, 3, or 10 mg/kg reconstituted in 5% DMAC in 0.5% methylcellulose and tumors were harvested 30 minutes after dosing. All tumors were processed for the determination of phospho-c-Met levels using the Human Phospho-HGFR/c-Met kit (R&D Systems). The plasma concentration of INCB28060 was determined by LC/MS/MS analysis following retro-orbital or cardiac puncture blood collection.

Results

INCB28060 is a potent and selective inhibitor of c-MET kinase

Through medicinal chemistry efforts, we have discovered a series of small molecule inhibitors of c-MET kinase, exemplified by INCB28060 (22). The enzymatic potency of INCB28060 was determined in an *in vitro* kinase assay using recombinant human c-MET and a biotin-labeled tyrosine-containing peptide as the substrate. INCB28060

Table 1. Enzymatic and cellular activities of INCB28060 against human c-MET

Assay	Enzyme or cell source	IC ₅₀ , ^a nmol/L	N
<i>In vitro</i> c-MET kinase assay	Recombinant human wild-type c-MET	0.13 ± 0.05	17
Cellular phospho-c-MET assay			
Constitutive	SNU-5 (gastric)	1.1 ± 0.4 ^c	28
	S114 (transfected)	0.9 ^c	2
HGF stimulated	A549 (lung)	0.7 ^c	2
	U-87MG (glioblastoma)	1 ^c	2
	786-0 (kidney)	0.6 ^c	2
	H441 (lung)	0.7 ^c	2
	H596 (lung) ^b	0.4 ^c	2
	H1437 (lung) ^b	0.3 ^c	2

^aValues are mean ± SD, and simple mean or estimated values for *n* = 2.

^bH596 and H1437 express mutant c-MET due to an exon 14 deletion and the R988C mutation, respectively (25).

^cValues were obtained using phospho-c-MET ELISA.

inhibits c-MET kinase activity with an average IC_{50} value of 0.13 nmol/L (Table 1). Additional kinetic studies suggest that INCB28060 is an ATP competitive and reversible inhibitor (data not shown). To determine the selectivity of INCB28060, the compound was profiled against a panel of 57 human kinases representing a structurally diverse cross-section of all known human kinases (Supplementary Table S1). At 2 μ mol/L, a concentration that is approximately 2,000-fold above the IC_{90} for c-MET, INCB28060 exhibited no more than 30% inhibition of all kinases tested, except for c-MET which was completely inhibited. Additional biochemical studies suggest that INCB28060 is inactive against $RON\beta$, another member of the c-MET RTK family, as well as EGFR and HER-3, members of the EGFR RTK family (Supplementary Fig. S1). Together, these data suggest that INCB28060 is a potent and highly selective c-MET kinase inhibitor.

INCB28060 effectively inhibits c-MET activation and signaling in cancer cells

The cellular activity of INCB28060 against human c-MET kinase was assessed by measuring inhibition of phospho-c-MET in relevant cancer cell lines where c-MET is activated. For example, the SNU-5 human gastric cancer cell line has constitutively activated c-MET as a result of *c-MET* amplification. As shown in Fig. 1A, INCB28060 inhibits c-MET phosphorylation with an IC_{50} value of approximately 1 nmol/L and a concentration of approximately 4 nmol/L inhibits c-MET more than 90%. The average IC_{50} value from multiple lots of the compound in this cell line was determined using a phospho-c-MET ELISA and is shown in Table 1. Similar inhibitory activity of INCB28060 was detected using other human cancer cell lines that represent tumor types where c-MET is frequently overexpressed or activated (Table 1). Noticeably, INCB28060 is slightly more potent in H596 and H1437 lung cancer cell lines that express mutant c-MET due to an exon 14 deletion and a R988C mutation, respectively. Both mutations are in the juxtamembrane domain of c-MET receptor (25) and perhaps adapt confirmations that are more favorable for compound binding. In addition, the activity of INCB28060 was assessed in the genetically engineered S114 mouse fibroblast cell line that expresses high levels of both human c-MET and HGF (26) and similar activity was observed. In addition, activity of INCB28060 against other c-MET phosphorylation sites such as tyrosine residues 1003 and 1349 was also assessed, and the compound exhibited similar potency in inhibiting phosphorylation of these residues (Supplementary Fig. S2). Inhibition of c-MET phosphorylation by INCB28060 treatment is reversible and the effect is significantly reduced in several hours after the compound is removed and completely disappeared by 48 hours (data not shown).

Given its strong activity against the c-MET enzyme, the potential effect of INCB28060 on c-MET-mediated signal transduction was further examined in cancer cells. INCB28060 effectively blocks phosphorylation of major downstream effectors of the c-MET pathway, including




Figure 1. INCB28060 inhibits human c-MET phosphorylation and c-MET-mediated signaling in the SNU-5 human gastric cancer cell line. A, SNU-5 cells were treated with INCB28060 at the indicated concentrations for 2 hours before harvesting and lysis. Equal amounts of protein extracts were subjected to Western blot analysis using anti-phospho-c-MET and anti-total c-MET antibodies. Inhibition of phospho-c-MET is shown in a representative Western blot analysis. B, SNU-5 cells treated with INCB28060 at the indicated concentrations for 2 hours were lysed and subjected to Western blot analyses using antibodies that recognize the individual (phospho-) proteins studied. A representative experiment shows that INCB28060 effectively inhibits phosphorylation of c-MET as well as c-MET pathway downstream effectors such as ERK1/2, AKT, FAK, GAB1, and STAT3/5.

ERK1/2, AKT, FAK, GAB1, STAT3/5 in SNU-5 cells (Fig. 1B). Strikingly, all the signaling proteins examined are effectively inhibited more than 90% by the compound at a concentration of approximately 4 nmol/L. Similar inhibitory effects were also observed in H1993 and A549 human lung cancer lines (data not shown). The data suggest that INCB28060, via inhibition of c-MET kinase activity, is capable of effectively blocking c-MET-mediated signaling cascades that are important for promoting cancer cell growth, survival, and invasion.

INCB28060 inhibits c-MET-dependent neoplastic phenotypes of cancer cells

Increased c-MET activity triggers a highly diverse set of signaling cascades, resulting in pleiotropic effects on tumor cells (1, 2). Therefore, INCB28060 was evaluated for its ability to block c-MET-dependent neoplastic activities in cancer cells, including cell proliferation (anchorage dependent and independent), migration, and antiapoptosis. As a result of *c-MET* amplification, the growth of SNU-5 cells is primarily driven by the activated c-MET kinase. INCB28060 inhibited SNU-5 viability or proliferation with an average




Figure 2. INCB28060 inhibits c-MET-dependent cell proliferation and survival. A, the indicated cell lines were cultured for 3 days in appropriate media containing 2% FBS in the presence of various concentrations of INCB28060, and cell viability was measured by MTS assay. The composite curve of percentage inhibition of cell viability by INCB28060 was plotted using average values of multiple experiments \pm SD. B, Western blot detection of c-MET protein in SNU-5 and SNU-1 cells. C, SNU-5 cells were grown in low-serum medium overnight and treated with INCB28060 at the indicated concentrations for 20 hours. DNA fragmentation analysis was carried out with the treated cells, and representative data showing levels of DNA fragmentation from 3 independent experiments are presented. D, SNU-5 cells receiving similar treatments to (C) were lysed for protein extraction and then subjected to Western blot detection of cleaved PARP. Result of a representative experiment is presented. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

IC₅₀ value of 1.2 nmol/L and a calculated IC₉₀ value of 4.6 nmol/L (Fig. 2A and Table 2). In contrast, the compound was inactive against the SNU-1 human gastric cancer cell line that does not express c-MET (Fig. 2A and B) and the HEK293 human embryonic kidney cell line, an immortalized nontumorigenic cell line that expresses low levels of c-MET (Table 2). When assessed against S114 cells, INCB28060 did inhibit cell proliferation, but with an average IC₅₀ value of 12.4 nmol/L (Table 2). The fact that the compound is less active against S114 cells versus SNU-5 cells is likely due to the unusually high levels of both c-MET and HGF expressed that make S114 cells more resistant to c-MET inhibition. Collectively, the data suggest that INCB28060 exhibits potent inhibitory activity only in cancer cells whose growth is specifically driven by the activated c-MET pathway.


The ability of INCB28060 to block anchorage-independent cancer cell growth was evaluated in a soft agar colony formation assay conducted with the U-87MG glioblastoma or H441 lung cancer cell line where the c-MET pathway is active (27). The compound is highly active with an estimated IC₅₀ value of approximately 2 nmol/L in U-87MG cells or 0.5 nmol/L in H441 cells (Fig. 3A and Table 2). Notably, there was almost no U-87MG cell growth upon the treatment of 16 nmol/L INCB28060. Another important function of the activated c-MET pathway is its ability to promote cell migration. To determine the effect of INCB28060 on cancer cell migration, a wound-healing-based migration assay was carried out with H441 cells. INCB28060 prevented HGF-stimulated H441 cell migration, with an IC₅₀ value of approximately 2 nmol/L (Fig. 3B and Table 2). Again, there was

Table 2. Activity of INCB28060 in cell-based functional assays

Assay	Cell line	IC ₅₀ , ^a nmol/L	N
Cell viability/proliferation	SNU-5	1.2 \pm 0.4	28
	S114	12.4	2
	SNU-1	>10,000	2
	HEK293	>20,000	2
Soft agar colony formation	H441	~0.5	2
U-87MG	~2	2	2
Migration/wound healing	H441	~2	2
Apoptosis/DNA fragmentation	SNU-5	Detectable at 1 nmol/L	3

^aValues are mean \pm SD, simple mean, or estimated values in case of semiquantitative assays for $n = 2$.

Figure 3. INCB28060 prevents anchorage-independent cancer cell growth and cell migration. Representative data of 2 independent experiments are presented to show the inhibitory effects of INCB28060 on anchorage-independent U-87MG cell growth in soft agar (A) and HGF-stimulated H441 cell migration (B). 28060, INCB28060.



little cell migration at a concentration of 16 nmol/L INCB28060.

Increased c-MET activity promotes cancer cell survival. As a consequence, cancer cells carrying activated c-MET are more resistant to apoptosis than cancer cells with a normal level of c-MET activity. To determine if INCB28060 treatment impacts c-MET-mediated cancer cell survival, DNA fragmentation, an indicator of cells undergoing apoptosis, was measured in SNU-5 cells treated with INCB28060. The compound effectively induced DNA fragmentation (Fig. 2C and Table 2). Furthermore, PARP cleavage, a biochemical read-out for apoptotic cells, was markedly induced by INCB28060 (Fig. 2D). A similar induction of PARP cleavage was readily detected in H441 cells upon the treatment of INCB28060 (data not shown). The results suggest that activated c-MET renders cancer cells resistant to apoptosis and INCB28060 can antagonize this activity.

INCB28060 shows strong antitumor activity in c-MET-dependent mouse tumor models

To assess the *in vivo* activities of INCB28060, we used the S114 cell-derived mouse tumor model. Because S114 cells express both human c-MET and HGF, tumors from these cells are dependent upon c-MET signaling for their growth. To determine the minimum dose of INCB28060 necessary to control c-MET phosphorylation, we orally administered to mice increasing doses of INCB28060 and measured phospho-c-MET levels in tumors 30 minutes later. As seen in Fig. 4A, 0.03 mg/kg INCB28060, the lowest dose tested, causes approximately 50% inhibition of c-MET phosphorylation. Escalating doses affect phospho-c-MET in a dose-dependent fashion, and single doses of 0.3 mg/kg or more resulted in greater than 90% inhibition. To further characterize the impact of INCB28060 over time, a single dose of 3 mg/kg was selected. Inhibition of phospho-c-MET exceeded 90% through the 7-hour measurement time point (Fig. 4B), which is consistent with the compound exposure exceeding protein-adjusted IC₉₀ (~71 nmol/L) for phospho-c-MET during the same period of time (Fig. 4B). There-

fore, the activity of INCB28060 is dose dependent and sustained over time as a result of effective drug exposure levels for that same period of time *in vivo*. Similar results were observed with the MKN-45 human gastric cancer cell-derived mouse tumor model that is driven by c-MET activation as a result of c-MET amplification (data not shown).

To determine how inhibition of c-MET phosphorylation translates to antitumor efficacy in the S114 model, tumor-bearing mice were treated orally with 3, 10, or 30 mg/kg twice daily. Dose-dependent inhibition of tumor growth was seen (Fig. 4C). Notably, complete regressions were seen in all mice in the group receiving 30 mg/kg INCB28060. Data obtained in the more sensitive U-87MG glioblastoma model, which expresses lower levels of endogenous c-MET and HGF, showed a similar dose-dependent inhibition of tumor growth with 35% and 76% tumor growth inhibition for groups dosed with 1 and 3 mg/kg INCB28060 once daily, respectively (Fig. 4D). Furthermore, once daily dosing of 10 mg/kg INCB28060 resulted in partial regressions in 6 of 10 U-87MG tumor-bearing mice. It is noted that in both S114 and U-87MG models, tumor growth inhibition increases with increased exposure of the compound and that tumor regressions could only be achieved when the compound exposure consistently exceeded 90% of c-MET inhibition (Fig. 4C). In these studies, INCB28060 was well tolerated at all doses during the treatment periods, with no evidence of overt toxicity or weight loss. Taken together, these data support the utility of INCB28060 in controlling the growth of c-MET driven tumors with maximal effects detected at IC₉₀ coverage.

c-MET positively regulates EGFR and HER-3 signaling pathways

Cross-talk between the c-MET pathway and other signaling proteins or pathways has been documented (10). To investigate the potential interactions between c-MET and EGFR family members such as EGFR and HER-3, we first determined if activated c-MET regulated expression of ligands for EGFR and HER-3 in human cancer cells.





Figure 4. INCB28060 inhibits c-MET phosphorylation and tumor growth in c-MET-dependent mouse tumor models. Equal amounts of protein extracts of tumor tissues from S114 tumor-bearing mice were analyzed for phospho-c-MET levels. Representative data are presented for dose-dependent inhibition of phospho-c-MET by INCB28060 at 30 minutes postdose (A) and for inhibition of phospho-c-MET over time by a single dose of 3 mg/kg INCB28060 (B). In (A) and (B), a group of 8 mice was used to generate each data value, and mean \pm SEM are shown. For INCB28060 exposure, mean plasma concentrations are presented. Representative data of dose-dependent tumor growth inhibition are shown in the S114 model (C) and in the U-87MG model (D). Mean tumor volumes \pm SEM are shown for the S114 model ($n = 8$ mice per group) and U-87MG model ($n = 10$ mice per group). For INCB28060 exposure, mean plasma concentrations \pm SEM for the first 16 hours of day 1 treatment are shown for the S114 model ($n = 8$ mice per group) in (C). PBAdjIC₉₀, protein binding-adjusted IC₉₀. The arrow-headed lines below the graphs in (C) and (D) indicate the days when the animals were treated with the compound.

Recombinant human HGF was used to induce c-MET activation in multiple human cancer cell lines including H596, A549, 786-0, H441, and H1993, and the levels of EGFR and HER-3 ligands such as TGF- α , amphiregulin (AR), and heregulin- β 1 (HRG- β 1) were measured in cell culture supernatants. As shown in Fig. 5A, induction of TGF- α , AR, and HRG- β 1 was detected in most of the cell lines tested. Interestingly, this induction could be effectively inhibited by INCB28060 treatment at potencies within the single-digit nanomolar range, as shown in H441 cells (Fig. 5B), suggesting that the induction of these ligands was indeed mediated through HGF-stimulated c-MET signaling. Furthermore, INCB28060 treatment also reduced unstimulated baseline levels of the ligands such as AR (Fig. 5C). Mechanistically, HGF/c-MET signaling could regulate the expression of these ligands via different mechanisms, including regulation of transcription and/or translation, or posttranslational regulation. To further explore the potential mechanism, we studied the effects of known inhibitors of RNA or protein synthesis on this regulatory process. We found that inclusion of either actinomycin D or cycloheximide at the concentrations previously shown to inhibit RNA or protein synthesis (28), significantly reduced the induction of

both AR and TGF- α by HGF (data not shown), suggesting that the regulation occurs at both transcriptional and translational levels.

In addition to the regulation of EGFR and HER-3 ligand expression, we further questioned if c-MET also regulates EGFR and HER-3 themselves. Despite the fact that INCB28060 is inactive against EGFR or HER-3 (Supplementary Fig. S1B and S1C), we found that in the c-MET amplified H1993 lung cancer cell line, INCB28060 treatment suppressed phosphorylation of both EGFR and HER-3 rapidly and as effectively as the compound inhibited c-MET phosphorylation (Fig. 6A). The inhibition of phosphorylated EGFR and HER-3 by INCB28060 persisted over a period of 24 hours (Fig. 6B). Because the H1993 cell line has constitutively active c-MET as a result of c-MET amplification, addition of exogenous HGF showed little effect on the phosphorylation status of either EGFR or HER-3 with or without INCB28060 pretreatment (Fig. 6B). Intriguingly, treatment of INCB28060 over a longer period (e.g., 24 hours) led to a dose-dependent increase in HER-3 protein levels, whereas EGFR protein levels were unchanged (Fig. 6B). Similar effects on total HER-3 and phospho-HER-3 (although to a lesser extent due to the excessive amount of total HER-3 induced by INCB28060 treatment for 24 hours)

Figure 5. HGF induces production of TGF- α , AR, and HRG- β 1 in cancer cells and INCB28060 effectively blocks the induction. **A**, the indicated human cancer cell lines were cultured in the presence or absence of 50 ng/mL recombinant human HGF for 24 hours and levels of TGF- α , AR, and HRG- β 1 in culture supernatants were measured by respective ELISAs. Representative data of 3 independent experiments are shown in the graphs. **B**, H441 cells were treated with 50 ng/mL HGF for 24 hours in the presence of various concentrations of INCB28060. The activity (IC_{50}) of the compound in inhibiting the induction of TGF- α and AR in supernatants was measured, and representative data of 3 independent experiments are shown. **C**, the indicated cell lines were cultured in the presence of various concentrations of INCB28060. Unstimulated baseline levels of AR were measured in culture supernatants 24 hours later and representative data are presented.



by INCB28060 treatment were also observed in the *c-MET* amplified SNU-5 cells, suggesting that in the *c-MET* amplified cancer cells, constitutively activated *c-MET* positively regulates the basal activity of both EGFR and HER-3. The fact that inhibition of *c-MET* kinase activity by INCB28060 induced HER-3 expression, however, suggests that *c-MET* may specifically suppress HER-3 expression in these cells. In addition to the regulation of activity and/or expression of EGFR and HER-3, direct physical associations between *c-MET* and EGFR or HER-3 were also detected in SNU-5 cells by immunoprecipitation (Fig. 6C and D). Similar immunoprecipitation experiments carried out using protein extract prepared with other human cancer cell lines showed very weak HER-3 detection after 24 hour compound treatment, likely because of low baseline HER-3 expression in those cell lines (Fig. 6D). Collectively, the data suggest that activated *c-MET* may regulate EGFR and HER-3 via multiple mechanisms, including stimulation of EGFR and HER-3 ligand production as well as activation of the receptors directly.

To further assess the relevance of HGF and *c-MET* in cancer patients, circulating HGF levels were measured in patients with various tumor types. HGF levels were significantly higher in the majority of patients with all tumor types tested (Supplementary Fig. S3). Approximately 49% of patients showed a 2-fold or greater increase than healthy volunteers. The data confirms that HGF levels are frequently elevated in cancer patients and might contribute to *c-MET* activation.

Discussion

The *c-MET* pathway is frequently dysregulated in a wide variety of human cancers and may play critical roles in cancer formation, progression, and dissemination, as well as resistance to approved therapies. Therefore, inhibition of increased *c-MET* signaling could have significant potential for the treatment of human cancers where the *c-MET* pathway is activated. In fact, recent clinical trials of *c-MET* pathway targeted agents have yielded convincing evidence to support the potential utility of this class of agents in treating various human cancers (29–34). However, there may be limitations for the *c-MET* pathway targeted agents that are in development (29). For example, most of the reported *c-MET* kinase inhibitors being evaluated in the clinic are not very selective for *c-MET* and exhibit activities against multiple kinases, which resulted in unwanted off-target toxicities (29, 32, 33). Furthermore, some of these inhibitors are not very potent and may have difficulty in achieving optimal *c-MET* inhibition in patients (29, 31, 35). On the other hand, several antibodies that recognize either human HGF or *c-MET* may have issues as large molecule biological agents in development (10). In addition, the antibodies specifically against HGF may have their activity limited to those patients where *c-MET* is activated by HGF, but not by many HGF-independent mechanisms (29, 34). Therefore, a truly selective and potent *c-MET* inhibitor with favorable pharmacologic properties could have multiple advantages.




Figure 6. Cross-talk between c-MET and EGFR or HER-3 in cancer cells. **A**, H1993 cells were treated with INCB28060 for 20 minutes at the indicated concentrations and whole-cell lysates were subjected to Western blot detections of phospho-c-MET or total c-MET, EGFR, and HER-3. Representative data of 3 independent experiments are presented. **B**, representative data of H1993 cells pretreated with dimethyl sulfoxide (DMSO) at a concentration equivalent to that in the treatment of 100 nmol/L INCB28060 or indicated concentrations of INCB28060 for 24 hours and followed by HGF stimulation for 15 minutes. **C**, SNU-5 cells were treated with 0.5 $\mu\text{mol/L}$ INCB28060 for 24 hours, and whole-cell lysates were immunoprecipitated (IP) with an anti-c-MET antibody, and total c-MET, EGFR, and HER-3 in the immunoprecipitates were detected by Western blotting with respective antibodies. **D**, representative data from similar analyses using various human cancer cell lines showing direct association between c-MET and HER-3 are presented. In addition, representative straight Western blot analyses of total and phospho-HER-3 using the cell lines treated with INCB28060 for 24 hours and straight Western blot analyses of total HER-3 using the cell lines treated with INCB28060 for 2 hours or 20 minutes are shown, respectively.

Here we describe the discovery and preclinical characterization of INCB28060, a novel selective c-MET kinase inhibitor that is currently being evaluated in phase I trials in advanced cancer patients (36). INCB28060 has been shown exceptionally potent activity against the c-MET enzyme and was shown to inhibit c-MET-mediated signaling and neoplastic activities in a variety of relevant human cancer cell lines, with low single-digit nmol/L potency. INCB28060 is highly specific for c-MET with greater than 10,000-fold selectivity over 56 other human kinases tested. In c-MET-dependent mouse tumor models, INCB28060 exhibited dose-dependent antitumor activity and caused tumor regression in some tumor-bearing mice at doses that exceeded IC_{90} coverage and the compound was well tolerated. Collectively, the data suggest that INCB28060 possesses potent *in vitro* and *in vivo* biological and pharmacologic activities, and further support its clinical development as a potentially effective oral treatment for human cancers. In addition, the remarkable biological and pharmacologic properties that INCB28060 has shown may allow this agent to overcome the aforementioned limitations of other c-MET pathway inhibitors and to specifically assess the true therapeutic potential of c-MET inhibition in cancer patients.

Another interesting finding from our characterization of INCB28060 is that inhibition of c-MET signaling might have additional benefit, as our data show that activated

c-MET also positively regulates other important cancer-promoting pathways such as EGFR and HER-3. Our study suggests that activated c-MET could upregulate the activity of EGFR and HER-3 through these receptors themselves and by increasing their ligand production, and that these effects could be effectively reversed by INCB28060. The data are generally consistent with previous observations (10, 28). To our surprise, however, we also found that activated c-MET specifically suppresses HER-3 expression but not EGFR expression in the c-MET amplified human cancer cell lines. This suppression of HER-3 expression by activated c-MET might be a common mechanism because an increased level of HER-3 protein physically associated with c-MET was also noted in other cell lines treated with INCB28060, prior to immunoprecipitation with a c-MET antibody (Fig. 6D). However, because HER-3 is expressed at fairly low levels in some tested cell lines and the HER-3 protein weakly detected by immunoprecipitation showed a slightly different size in 1 or 2 tested cell lines, further study of this issue using other model cell lines will be helpful. The fact that the induction of HER-3 protein could only be detected after longer periods of compound treatment indicates that the effect may be regulated at the transcriptional or posttranscriptional level. Furthermore, our data can not rule out the possibility that c-MET itself dose not directly suppress HER-3 expression. Instead,

c-MET may regulate other intracellular proteins or enzymes that in turn suppress HER-3 expression. Additional studies are necessary to further elucidate the potential mechanism by which c-MET downregulates HER-3 expression in these cell lines.

Despite the observation that the expression of HER-3 is suppressed by activated c-MET, the basal activity of HER-3 is upregulated by c-MET in *c-MET* amplified cancer cells, as INCB28060 treatment led to a strong inhibition of phosphorylated HER-3. Thus, the net effect of c-MET inhibition on HER-3 activity is still inhibitory in these cells. Because development of drug resistance is common in cancer patients and cross-talk between different cancer-promoting pathways might be one of the mechanisms that confers drug resistance, the inhibitory effects of a

c-MET inhibitor like INCB28060 on c-MET signaling as well as pathways like EGFR and HER-3 that are regulated by c-MET may help to reduce drug resistance in some patients. Future clinical studies with INCB28060 may shed light on this issue.

Disclosure of Potential Conflicts of Interest

All authors are employees of and/or own stock in Incyte Corporation.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received May 2, 2011; revised September 1, 2011; accepted September 8, 2011; published OnlineFirst September 14, 2011.

References

- Birchmeier C, Birchmeier B, Gherardi E, Vande Woude GF. Met, metastasis, motility and more. *Nat Rev Mol Cell Biol* 2003;4:915–25.
- Christensen JG, Burrows J, Salgia R. c-Met as a target for human cancer and characterization of inhibitors for therapeutic intervention. *Cancer Lett* 2005;225:1–26.
- Roos F, Ryan AM, Chamow SM, Bennett GL, Schwall RH. Induction of liver growth in normal mice by infusion of hepatocyte growth factor/scatter factor. *Am J Physiol* 1995;268:G380–6.
- Michalopoulos GK, DeFrances MC. Liver regeneration. *Science* 1997;276:60–6.
- Matsumoto K, Nakamura T. Hepatocyte growth factor: renotropic role and potential therapeutics for renal diseases. *Kidney Int* 2001;59:2023–38.
- Nakamura T, Mizuno S, Matsumoto K, Sawa Y, Matsuda H, Nakamura T. Myocardial protection from ischemia/reperfusion injury by endogenous and exogenous HGF. *J Clin Invest* 2000;106:1511–9.
- Rabkin R, Fervenza F, Tsao T, Sibley R, Friedlaender M, Hsu F, et al. Hepatocyte growth factor receptor in acute tubular necrosis. *J Am Soc Nephrol* 2001;12:531–40.
- Van Andel Institute. Hepatocyte growth factor/scatter factor, Met and cancer references [database on the Internet]. Grand Rapids (MI): Van Andel Institute. Available from: <http://www.vai.org/met/>.
- Schmidt L, Junker K, Nakaigawa N, Kinjerski T, Weirich G, Miller M, et al. Novel mutations of the MET proto-oncogene in papillary renal carcinomas. *Oncogene* 1999;18:2343–50.
- Liu X, Yao W, Newton RC, Scherle PA. Targeting the c-MET signaling pathway for cancer therapy. *Expert Opin Invest Drugs* 2008;17:999–1011.
- Zeng ZS, Weiser MR, Kuntz E, Chen CT, Khan SA, Forslund A, et al. c-MET gene amplification is associated with advanced stage colorectal cancer and liver metastases. *Cancer Lett* 2008;265:258–69.
- Galeazzi E, Olivero M, Gervasio FC, De Stefani A, Valente G, Comoglio PM, et al. Detection of MET oncogene/hepatocyte growth factor receptor in lymph node metastases from head and neck squamous cell carcinomas. *Eur Arch Otorhinolaryngol* 1997;254 Suppl 1: S138–43.
- Graveel C, Su Y, Koeman J, Wang LM, Tessarollo L, Fiscella M, et al. Activating Met mutations produce unique tumor profiles in mice with selective duplication of the mutant allele. *Proc Natl Acad Sci U S A* 2004;101:17198–203.
- Ponzo MG, Lesurf R, Petkiewicz S, O'Malley FP, Pinnaduwa D, Andrulis IL, et al. Met induces mammary tumors with diverse histologies and is associated with poor outcome and human basal breast cancer. *Proc Natl Acad Sci U S A* 2009;106:12903–8.
- Seiwert TY, Jagadeeswaran R, Faoro L, Janamanchi V, Nallasura V, El Dinali M, et al. The MET receptor tyrosine kinase is a potential novel therapeutic target for head and neck squamous cell carcinoma. *Cancer Res* 2009;69:3021–31.
- Knowles LM, Stabile LP, Eglhoff AM, Rothstein ME, Thomas SM, Gubish CT, et al. HGF and c-Met participate in paracrine tumorigenic pathways in head and neck squamous cell cancer. *Clin Cancer Res* 2009;15:3740–50.
- Okuda K, Sasaki H, Yukiue H, Yano M, Fujii Y. Met gene copy number predicts the prognosis for completely resected non-small cell lung cancer. *Cancer Sci* 2008;99:2280–5.
- Toiyama Y, Miki C, Inoue Y, Okugawa Y, Tanaka K, Kusunoki M, et al. Serum hepatocyte growth factor as a prognostic marker for stage II or III colorectal cancer patients. *Int J Cancer* 2009;125:1657–62.
- Gupta A, Karakiewicz PI, Roehrborn CG, Lotan Y, Zlotta AR, Shariat SF, et al. Predictive value of plasma hepatocyte growth factor/scatter factor levels in patients with clinically localized prostate cancer. *Clin Cancer Res* 2008;14:7385–90.
- Cappuzzo F, Marchetti A, Skokan M, Rossi E, Gajopathy S, Felicioni L, et al. Increased MET gene copy number negatively affects survival of surgically resected non-small cell lung cancer patients. *J Clin Oncol* 2009;27:1667–74.
- Kammula US, Kuntz EJ, Francone TD, Zeng Z, Shia J, Landmann RG, et al. Molecular co-expression of the c-Met oncogene and hepatocyte growth factor in primary colon cancer predicts tumor stage and clinical outcome. *Cancer Lett* 2007;248:219–28.
- Weng L, Qiao L, Zhou J, Liu P, Pan Y, inventors; Incyte Corporation, assignee. Salts of 2-fluoro-N-methyl-4-[7-(quinolin-6-yl-methyl)-imidazo[1,2-b][1,2,4]triazin-2-yl]benzamide and processes related to preparing the same. Europe patent WO 2009143211. 2009 Nov 26.
- Behshad E, Klabe RM, Margulis A, Becker-Pasha M, Rupar MJ, Collier P, et al. Phosphorylation state-dependent high throughput screening of the c-MET kinase. *Curr Chem Genomics* 2010;4:27–33.
- Liu X, Fridman JS, Wang Q, Caulder E, Yang G, Covington M, et al. Selective inhibition of ADAM metalloproteases blocks HER-2 extracellular domain (ECD) cleavage and potentiates the anti-tumor effect of trastuzumab. *Cancer Bio Ther* 2006;5:648–56.
- Kong-Beltran K, Seshagiri S, Zha J, Zhu W, Bhawe K, Mendoza N, et al. Somatic mutations lead to an oncogenic deletion of met in lung cancer. *Cancer Res* 2006;66:283–9.
- Rong S, Bodescot M, Blair D, Dunn J, Nakamura T, Mizuno K, et al. Tumorigenicity of the met proto-oncogene and the gene for hepatocyte growth factor. *Mol Cell Biol* 1992;12:5152–8.
- Zou HY, Li Q, Lee JH, Arango ME, McDonnell SR, Yamazaki S, et al. An orally available small-molecule inhibitor of c-MET, PF-2341066, exhibits cytoreductive antitumor efficacy through antiproliferative and antiangiogenic mechanisms. *Cancer Res* 2007;67:4408–17.
- Reznik TE, Sang Y, Ma Y, Abounader R, Rosen EM, Xia S, et al. Transcription-dependent epidermal growth factor receptor activation by hepatocyte growth factor. *Mol Cancer Res* 2008;6: 139–50.

29. Liu X, Newton RC, Scherle PA. Developing c-MET pathway inhibitors for cancer therapy: progress and challenges. *Trends Mol Med* 2010;16:37–45.
30. Spigel D, Ervin T, Ramlau R, Daniel D, Goldschmidt J, Krzakowski M, et al. Randomized multicenter double-blind placebo-controlled phase II study evaluating MetMAb, an antibody to Met receptor, in combination with erlotinib, in patients with advanced non-small cell lung cancer [abstract]. In: The 35th Congress of the European Society for Medical Oncology; 2010. Abstract number 5146.
31. Mekhail T, Rich T, Rosen L, Chai F, Semic-Suka Z, Savage RE, et al. Final results: a dose escalation phase I study of ARQ197, a selective c-Met inhibitor, in patients with metastatic solid tumors. *J Clin Oncol* 27:15s, 2009; (suppl; abstr 3548).
32. Eder JP, Shapiro GI, Appleman LJ, Zhu AX, Miles D, Keer H, et al. A phase I study of foretinib, a multi-targeted inhibitor of c-Met and vascular endothelial growth factor receptor 2. *Clin Cancer Res* 2010;16:3507–16.
33. Kurzrock R, Sherman S, Hong D, Ng C, Frye J, Janisch L, et al. A phase I study of XL184, a MET, VEGFRs, and RET kinase inhibitor, administered orally to patients (pts) with advanced malignancies, including a subgroup of pts with medullary thyroid cancer (MTC) [abstract]. In: The AACR-NCI-EORTC International Conference, Molecular Targets and Cancer Therapeutics, Discovery, Biology, and Clinical Applications. 2008 Oct 21–24; Geneva, Switzerland. Philadelphia (PA): AACR; 2008. Abstract nr 379.
34. Gordon MS, Sweeney CS, Mendelson DS, Eckhardt SG, Anderson A, Beaupre DM, et al. Safety, pharmacokinetics, and pharmacodynamics of AMG 102, a fully human hepatocyte growth factor-neutralizing monoclonal antibody, in a first-in-human study of patients with advanced solid tumors. *Clin Cancer Res* 2010;16:699–710.
35. Munshi N, Jeay S, Li Y, Chen CR, France DS, Ashwell MA, et al. ARQ 197, a novel and selective inhibitor of the human c-Met receptor tyrosine kinase with antitumor activity. *Mol Cancer Ther* 2010;9:1544–53.
36. Clinicaltrials.gov. A phase I, open-label, dose-escalation study to determine the safety, tolerability, pharmacokinetics, and pharmacodynamics of INCB028060 in subjects with advanced malignancies. Bethesda, MD: NIH. Available from: <http://clinicaltrials.gov/ct2/results?term=INCB028060>.